

Original citation:

Karasouli, E., et al. (2011). All-cause mortality after non-fatal self-poisoning : a cohort study. *Social Psychiatry and Psychiatric Epidemiology*, 46(6), pp. 455-462.

Permanent WRAP url:

<http://wrap.warwick.ac.uk/37273>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work by researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Publisher's statement:

The final publication is available at link.springer.com

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk

warwick**publications**wrap

highlight your research

<http://go.warwick.ac.uk/lib-publications>

Manuscript Number: SPPE-D-09-00261R1

Title: All-cause mortality after non-fatal self-poisoning: a cohort study

Article Type: Original Paper

Keywords: self-harm; self-poisoning; mortality; natural causes of death; physical illness

Corresponding Author: Miss Eleni Karasouli,

Corresponding Author's Institution: University of Leeds

First Author: Eleni Karasouli

Order of Authors: Eleni Karasouli; David Owens; Rachel L Abbott; Keith M Hurst; Michael Dennis

Abstract: Background. Suicide has been repeatedly shown to have greatly increased incidence after non-fatal self-harm but far less is known about early death from other causes. The present study's aim was to describe mortality and risk factors concerning all causes of death after non-fatal self-poisoning. Methods. A prospective cohort study of 976 patients who attended the Emergency Department in Nottingham, UK during a nine-month period in 1985-86. Information on deaths was obtained for 16 years following an episode of self-poisoning, from the records of the Office for National Statistics. Results. The observed:expected ratio for all-cause mortality was 2.2. Deaths due to diseases of the digestive and respiratory systems were respectively 4.4 and 2.9 times more frequent than expected. The risk for accidents was 6-fold and for probable suicides 17-fold compared to the risk in the general population. The main risk factor for subsequent deaths from natural causes was increasing age. Conclusions. The findings of this study suggest that patients who survive self-poisoning have an increased risk of death from natural and unnatural causes. The findings point towards the need for more effective clinical management and preventive initiatives.

ABSTRACT

Background. Suicide has been repeatedly shown to have greatly increased incidence after non-fatal self-harm but far less is known about early death from other causes. The present study's aim was to describe mortality and risk factors concerning all causes of death after non-fatal self-poisoning.

Methods. A prospective cohort study of 976 patients who attended the Emergency Department in Nottingham, UK during a nine-month period in 1985-86. Information on deaths was obtained for 16 years following an episode of self-poisoning, from the records of the Office for National Statistics.

Results. The observed:expected ratio for all-cause mortality was 2.2. Deaths due to diseases of the digestive and respiratory systems were respectively 4.4 and 2.9 times more frequent than expected. The risk for accidents was 6-fold and for probable suicides 17-fold compared to the risk in the general population. The main risk factor for subsequent deaths from natural causes was increasing age.

Conclusions. The findings of this study suggest that patients who survive self-poisoning have an increased risk of death from natural and unnatural causes. The findings point towards the need for more effective clinical management and preventive initiatives.

Keywords:

Self-harm, self-poisoning, mortality, natural cause of death, physical illness, risk factors.

INTRODUCTION

Self-harm is an important issue for healthcare services. Rates of non-fatal self-harm are around their highest ever level with an estimate of 220,000 episodes dealt with by general hospitals in the UK each year [13]. Self-harm is one of the commonest reasons for medical admission to hospital, the commonest reason for females and the second commonest for males. Self-harm, especially when repetitive, is responsible for substantial health service costs [28]. Of all self-harm episodes resulting in hospital attendance, over 80% involve self-poisoning [13].

Many studies have focused on the association between self-harm and subsequent suicide (for example, [18, 27]), revealing a high rate of suicide after self-harm that has been estimated to be more than 60 times the general population rate [15]. The estimated rate of death by suicide after self-harm varies between studies, depending mainly on differing patient populations and follow-up times, but it is high; a systematic review estimated that more than 6 per cent of patients attending hospital due to self-harm will have died by suicide after ten years [23].

Although there have been numerous research reports that examine the relation between self-harm and subsequent suicide, much less attention has been paid to the risk of death from causes other than suicide. There is evidence to suggest that physical illness is associated with excess risk of suicide [11, 30] and attempted suicide [5]. A 10-year follow-up study in Denmark investigated mortality for patients admitted to hospital after self-poisoning [21]; many more deaths than expected were due to natural causes comprising respiratory, neurological, digestive, neoplasm and alcohol-related conditions. A study in Oslo followed up patients treated for self-poisoning for a 10-year period and

1 found suicide, heart disease, opiate abuse, and accidents to be the
2 commonest causes of death [9]. A systematic review and meta-analysis found
3 a raised incidence of early death from natural causes [19]. Similar results were
4 found more recently in follow-up studies conducted in Austria, Australia and
5 Finland among patients who had attended hospital because of self-poisoning
6 or attempted suicide; in each case there were clear excesses of deaths due to
7 natural and unnatural causes [1, 2, 22]. On the other hand, a Canadian study
8 found that patients admitted to inpatient psychiatric care following a suicide
9 attempt were significantly less likely than the population in general to die of
10 natural causes [17].
11
12
13
14
15
16
17
18
19
20
21
22
23

24 The only comparable UK study to have considered the relation between self-
25 harm and subsequent death due to reasons other than suicide is from Oxford
26 [16]. The study sample comprised all patients attending the general hospital in
27 Oxford over a period of 20 years from 1978 to 1997, following them up for an
28 average of 13 years – to the end of 2000. The study observed a higher than
29 expected death rate from natural causes, particularly from respiratory,
30 circulatory, neurological, endocrine, digestive, skin and musculoskeletal
31 conditions. Deaths due to accidents were also more frequent than expected,
32 and homicides were more frequent among males. The Oxford study is
33 particularly interesting because it used larger sample sizes and longer follow-
34 up periods than did the previous research cited.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Our investigation set out to corroborate or refute earlier findings about excess
52 mortality following non-fatal episodes of self-harm by following-up for 16 years
53 a rather different cohort: one that was assembled in a very short inception
54 period (less than one year), and from a large city that is more typical of urban
55 England. The aims of the present study are: 1) to determine the subsequent
56
57
58
59
60
61
62
63
64
65

mortality rates and causes of death in patients who had previously attended the Emergency Department because of non-fatal self-poisoning, 2) to examine the risk factors for subsequent non-suicidal deaths, and 3) to determine any possible differences between those who died by natural causes and those who were alive in the follow-up period. Findings on the increased risk of suicide in these patients have previously been published [25].

METHOD

Study population and sample

The cohort comprises 976 consecutive patients aged 14 years or over who attended a large Emergency Department in Nottingham as a consequence of self-poisoning over nine months during 1985-1986. The Emergency Department was the only one serving the city and adjacent population of over 600,000. Nottingham is a large city, its surrounding urban area being the seventh largest in the UK, with a socio-demographic profile typical of UK metropolitan areas. The index self-poisoning episode was defined as the first attendance at the Emergency Department because of self-poisoning during the study period from November 1985 to July 1986. National data on population and cause-specific mortality were obtained from the Office for National Statistics.

Definition of self-harm and self-poisoning

For the purpose of this report the definition of self-harm is intentional self-poisoning or self-injury, irrespective of motivation or degree of suicidal intent [15]. Self-poisoning is defined as the intentional ingestion of more than the prescribed amount of any drug, whether or not there is evidence that the act

was intended to result in death. The definition also includes intentional poisoning with non-ingestible substances and intentional overdoses of recreational drugs [13].

Patient Characteristics

The study patients were identified round the clock by Emergency Department clerical staff, and the clinical staff in the Emergency Department collected information on a research checklist while the patients were in the emergency room. Checklists asked about potential risk factors for the seriousness of the index episode as a suicide attempt and for the likelihood of adverse outcome in the medium to long term. Items included past mental health service contact, previous self-harm, living arrangements, and social and medical status. In addition data were collected about demographic variables, and about alcohol consumption around the time of the self-poisoning [24, 25].

Follow-up

The identifying particulars of patients from the 1985-1986 cohort were passed on to the Office for National Statistics (ONS) who ascertained whether the people were alive or dead at the end of 2002, some 16-17 years after attendance at the Emergency Department because of self-harm [25]. In the case of those who were deceased, ONS provided a copy of the death certificate, which includes the ICD-10 coding for the causes of death [32].

After careful examination of all death certificates it was decided that only the primary causes of death recorded would be used for this study. Secondary causes were either further descriptions of primary causes or they were not related to death. The natural causes of death were grouped as follows: neoplasms (ICD-10: C00-D48), diseases of the circulatory system (I00-I99),

diseases of the respiratory system (J00-J99), diseases of the digestive system (K00-K93), diseases of the genitourinary system (N00-N99), and deaths from all other causes. The unnatural causes of death were: accidents (S00-T98, V01-X59, Y40-Y98) and probable suicides (X60-X84, Y10-Y43). The category of probable suicides includes deaths recorded by coroners as suicides (X60-X84) or open verdicts (Y10-Y43). It is standard practice to combine open verdicts with suicides, as it has been suggested that suicide mortality is underestimated when suicide is used alone [4].

Statistical Analyses

Mortality rates, for all causes and for cause-specific deaths in the study population compared to the general population, were calculated using mid-year populations for England and Wales. The total of person-years at risk were calculated according to gender and 10-year age bands, with the exception of the last two intervals which ranged from 45 to 59 years and 60 and over – because of the right skewed age pattern seen in self-poisoning. Individuals aged 14 years and under were excluded due to age-group inconsistencies of the Office for National Statistics data. The expected number of deaths was obtained by multiplying the person-years at risk by the national rate.

Standardised mortality ratios (SMRs) were calculated by dividing the observed numbers of death by the expected numbers.

For the comparison of patients who were alive with those who had died, non-parametric two-tailed tests (Mann Whitney U test) were used in comparisons of metric variables, and two-tailed chi-squared tests when comparing proportions. We looked for predictors of subsequent death, analyzing the variables collected from Emergency Department records and the clinicians' checklists.

Survival analysis was conducted to create a Kaplan-Meier survival graph for assessing the survival time for each gender.

Using the variables that proved to have a statistically significant association with subsequent mortality, Cox's proportional hazards regression models were used to determine any independent predictors of death due to natural causes. Cases with missing variables were excluded. When Cox regression could not include all possible risk factors, because of insufficient outcome events, logistic regression analyses were used, in which case the final model identifies any additional variables but it does not take into account the time of follow-up. For selecting the model all variables were entered in one block, using the Enter method of regression. Hazard ratios and odds ratios are reported with 95% confidence intervals (95% CI). Goodness of fit for the final models was acceptable ($P < 0.001$). The data were analysed using SPSS version 16.0.

Ethical Approval

Ethical approval was received from the Faculty Research Ethics Committee (Faculty of Development and Society) of Sheffield Hallam University and the Nottingham (National Health Service) Research Ethics Committee.

RESULTS

Study sample

During the nine months of recruitment (1985-1986), 976 people attended the Emergency Department as a result of 1091 episodes of self-poisoning. 591 (61%) of those were female and 384 (39%) were male. The median age of the sample was 27 years, range 14 to 86, interquartile range 20 to 39 years.

Young adults made up most of the sample, with 84% ($n = 815$) under the age of 45 years.

Mortality

The Office for National Statistics was able to trace 912 (93%) of the 976 people. Individuals not traced were similar to those traced in terms of age and gender. 157 out of 912 people (17%) had died by the end of the follow-up period (at least 16%, 157/976, of the original cohort); 84 of the 591 women (14%), and 73 of the 384 men (19%). From those traced, 125 died by causes other than probable suicide (80% of the deaths). The median time to death, following the index episode of self-poisoning, for those who died by natural causes or accidents was 9.8 years for males ($n = 52$) and 7.2 years for females ($n = 73$). Survival analysis (Figure 1) shows that the risk of death is similar for each gender (logrank test, $\chi^2 = 0.17$, $P = 0.68$; Hazard ratio 0.92, 95% CI 0.62 to 1.37).

Figure 1 about here

Causes of death

The causes of death, by gender are shown in Table 1. By the end of 2002, 99 people had died from natural causes, 17 died by accidents, 32 died from probable suicides, and in 9 cases the cause was unknown or unspecified but we found no indications that the death was due to probable suicide. In the natural causes of death, the standardized mortality ratio (SMR) was highest for digestive diseases, at ratios between four- and five-fold for each gender. A high observed to expected ratio was also found for respiratory diseases, with ratios around 3. Very high mortality was observed for accidents, with female

SMR 7.6, and male SMR 5.3. As expected, from the findings reported previously about suicides [25], the SMR for suicide pointed towards a huge excess of probable suicides following non-fatal self-poisoning: in females SMR around 20 and in males 16.

Age-specific mortality

Table 1 also shows mortality ratios from all causes across 5 age bands. The mortality ratio from all causes was 1.96 (95% CI 1.5-2.4) for females and 2.43 (95% CI 1.9-3.0) for males. In the females, the excess of deaths was greatest (around 9-fold) between 25 and 34 years of age, for the males aged between 35 and 44 years (around 8-fold).

Table 1 about here

Patients who died by causes other than suicide, compared with patients who were alive

A number of differences were observed between patients who died either by natural cause or accident (that is, non-suicidal deaths) and patients who were alive at the end of follow-up. Differences that are not statistically significant are not reported here. As expected, patients who died in the follow-up period were significantly older at the time of index attendance than were patients who remained alive: median age at index episode of those who were alive at follow-up was 24 years (interquartile range 19 to 34 years), while median age at index of those who died during the follow-up period was 59 years (48 to 71), ($U = 11879$, $P < 0.001$).

Patients who had died showed an excess of a number of characteristics (Table 2). At the point of the index episode those who died but not by suicide were

much more likely than those who remained alive to be suffering from serious physical illness, and to be living alone – both factors that are closely associated with increasing age. They were also much more likely to have had psychiatric care, admission to a psychiatric unit, and previous self-harm. At the time of the episode of index attendance at the emergency department, those who subsequently died from causes other than suicide were more likely to have attended hospital during the working day, and to have ingested psychotropic drugs (rather than analgesics and a range of other medications) or non-ingestible substances. They were also much more likely to be drowsy or unconscious. It is also clear that there was a higher than expected non-suicidal mortality among patients who went on to repeat self-harm in the year following the index episode.

Table 2 about here

All of the above findings held true whether deaths designated as due to accidents were included or not with deaths from natural causes (Table 2).

Risk Factors for deaths from natural causes

In the regression models, the outcome was death from natural causes; we did not create a model for the outcome of natural and accidental deaths combined. A number of relations between single variables and mortality were observed, although Cox's proportional hazards regression model showed only one variable to have a clear effect when the factors were considered together: increasing age at the time of attendance because of self-harm (hazard ratio = 1.1 per year, 95% CI 1.09 -1.12).

The data were re-analysed using logistic regression in order to identify any other important variables. Again, only increasing age was found to independently predict death by natural causes (odds ratio = 1.13 per year, 95% CI 1.10 -1.16). The effects of the other variables that were significant with univariate analysis (Table 2) were not confirmed in the regression model.

DISCUSSION

This prospective cohort study, with its sample consisting of mainly young adults, showed high mortality with 17% of them dead by 16 years later. Overall, mortality was double the expected rate found in the general population. Females between the ages of 25 and 34 years had the highest excess of non-suicidal deaths with a standardised mortality ratio of about 9, while women aged 45 to 59 years had a standardised mortality ratio of over 5. The risk pattern was slightly different for males, whose highest excess of deaths was at 35 to 44 years (more than 7 times that of the general population), with a high risk also between the ages of 25 and 34 (standardised mortality ratio around 6).

Around 80% of all deaths were from natural causes or accidents, with most of the males having died within ten years and most of females by seven years. The overall risk of death during follow-up was similar for both genders.

As expected, the mortality was very high from probable suicide – around 16 to 20 times the expected rate. The mortality rate was, however, higher than expected even after the deaths by suicide were set aside. Excess risk of death was highest for accidents, but deaths from a range of natural causes were also substantially more frequent than expected in the general population. For

1 example, people who self-poisoned had more than four times the general
2 population risk of dying from digestive diseases and they were around three
3
4 times as likely to die from diseases of the respiratory system.
5
6
7

8 The people who died by natural causes or accidents showed a number of
9 differences from those who were alive at the end of follow-up. First, as
10
11 expected, they were older at the time of attendance at the Emergency
12
13 Department than were the sample of people who were still alive 16 years later.
14
15 They were also more likely to be living alone and to be suffering from a serious
16
17 illness at the time of overdose. Those who had died also showed strong
18
19 associations with previous and subsequent self-harm, and with past psychiatric
20
21 history. At the time of the index attendance, people who later died were more
22
23 often showing impaired consciousness, were more likely to have attended the
24
25 emergency department during the working day, and more likely to have
26
27 ingested psychotropic drugs or non-ingestibles.
28
29
30
31
32
33
34

35 The present study has a number of strengths. First, the study sample included
36
37 all patients attending the emergency department because of self-poisoning.
38
39 Many studies of outcome after self-harm have a filter based on physical or
40
41 psychiatric severity; for example, samples are drawn only from patients
42
43 assessed by the mental health team, or are even restricted to those admitted
44
45 to the mental health unit. Second, the use of national rather than local mortality
46
47 data resulted in high ascertainment of survival or death, with only 6% untraced.
48
49 Third, Nottingham has a socio-demographic profile typical of UK metropolitan
50
51 areas and the study findings are thereby likely to be relevant to other places. A
52
53 limitation of this study is that not all possible associations have been examined
54
55 due to few deaths in some categories of cause of death. Sample size also
56
57 meant that we were largely unable to identify independently predictive
58
59
60
61
62
63
64
65

variables - partly due to inconsistent recording of data by the emergency department staff at the time of the original hospital attendance for self-poisoning.

Our findings are broadly in line with previous research. High overall mortality after self-harm has been reported by other studies [8, 21, 10, 31]. The importance of physical illness at the time of self-harm was also evident in a cohort of young people [14]. Our study further shows that physical illness is a risk factor for subsequent death among people who have previously overdosed. Unfortunately, we do not have the data needed to examine whether the illness present when individuals self-harmed is similar to the type of illness which caused subsequent death.

An excess of deaths due to natural causes has been found in previous studies of self-harm [14, 26]. In particular, our findings are closely congruent with those from a large cohort study based on 20 years of episodes of non-fatal self-harm in Oxford, which found more than the expected number of deaths due to respiratory diseases, digestive disorders, and accidents [16]. An excess of digestive causes of death was also found in a Danish study [21]. Respiratory disorders are associated with long-term disability and it may well be the presence of disability in those with respiratory disorders that links with self-harm [7]. Our method of data collection from emergency department records meant that we were unable to determine whether people who died from digestive or respiratory diseases had self-poisoned with medicines used in their treatment for these disorders.

The significant excess of accidental deaths points to a number of possible interpretations including excessive risk-taking behaviour and the

1 consequences of mental disorder and associated treatment; but there is also a
2 possible misclassification of deaths that represent probable suicide. In the UK
3
4 and elsewhere, studies that have scrutinised certification of deaths with a
5
6 verdict of misadventure or accident found some that were deemed probable
7
8 suicides [6, 3]. A small number of suicides may be missed because the death
9
10 was not subject to a coroner's investigation; researchers in Sheffield, UK
11
12 examined clinicians' ability to recognise reportable deaths using fictitious case
13
14 histories and some clinicians failed to recognize probable suicides [29].
15
16
17
18
19

20 Emergency department records are ideal for ensuring that all patients
21
22 attending the hospital, not just those admitted or subjected to specialist
23
24 consultations, are included in analyses. These case records, however, typically
25
26 contain little information about patients' physical and mental health beyond that
27
28 needed for the management of the acute episode of poisoning. Consequently,
29
30 it is not possible to make many robust assertions from these data about why
31
32 mortality – from suicide, accident and natural causes – is so much higher
33
34 among people who have self-poisoned than in the general population. In
35
36 particular, neither the present study nor other published work sheds much light
37
38 on the reasons for an excess of deaths from natural causes. Our multivariate
39
40 analysis points towards no useful factors that can be modified either by
41
42 clinicians or through public health initiatives. Even the single variable analyses
43
44 show only relations between early death and background factors that cannot
45
46 realistically be modified, such as living alone, and having a history of physical
47
48 or psychiatric illness. All the features of the non-fatal episode of self-poisoning
49
50 that emerged as relevant – ingesting non-ingestible substances or prescribed
51
52 psychotropic medicines, attending during the day-time, and being rendered
53
54 drowsy or unconscious – are likely to be largely explained by age.
55
56
57
58
59
60
61
62
63
64
65

1 The current context for care and aftercare following hospital attendance due to
2 self-harm is of uncertainty about how best to deliver the most effective service.
3
4 The Cochrane review of interventions is inconclusive [12] and NICE guidelines
5
6 in England are based entirely on good practice points because the clinical
7
8 evidence from studies of good quality is unavailable [20]. People attending
9
10 hospital because of self-harm, more than a quarter of a million each year in the
11
12 UK, have a poor outcome as judged by non-fatal repetition, suicide, accident
13
14 and natural causes. The present study is an indication of how important it is to
15
16 move forwards towards an evidence based approach to assessment, treatment
17
18 and follow-up.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Fig. 1 Death from any cause according to gender: Kaplan-Meier curves represent time from the first episode during study period to death.




Table 1 Cause-specific and age-specific mortality ratios, by gender.

	Females			Males			Both	
	Number who died (N=84)	SMR	(95% CI)	Number who died (N=73)	SMR	(95% CI)	SMR	(95% CI)
Cause of Death								
Neoplasm	12	1.15	0.6-1.9	8	1.05	0.5-2.0	1.11	0.7-1.7
Circulatory	20	1.17	0.7-1.8	14	1.22	0.7-2.0	1.19	0.8-1.6
Respiratory	18	2.69	1.6-4.1	13	3.24	1.8-5.4	2.90	2.0-4.1
Digestive	7	4.16	1.8-8.2	5	4.91	1.8-10.9	4.44	2.4-7.5
Genitourinary	1	1.85	0.1-9.1	1	3.36	0.2-16.5	2.38	0.4-7.9
Accidents	8	7.60	3.5-14.4	9	5.32	2.6-9.8	6.19	3.7-9.7
Probable suicides	11	19.82	10.4-34.4	21	15.84	10.1-23.8	17.01	11.8-23.7
All ages	84	1.96	1.5-2.4	73	2.43	1.9-3.0	2.15	1.8-2.5
15-24	4	2.97	0.9-7.1	2	1.13	0.1-3.7	1.93	0.7-4.0
25-34	8	8.97	4.1-17.0	11	5.71	3.0-9.9	6.74	4.1-10.3
35-44	4	2.45	0.7-5.9	15	7.61	4.4-12.2	5.28	3.2-8.0
45-59	17	5.40	3.2-8.4	15	4.93	2.8-7.9	5.17	3.6-7.2
60+	51	1.42	1.0-1.8	30	1.40	0.9-1.9	1.41	1.1-1.7

SMR: Standardised Mortality Ratio; CI: Confidence Interval

Table 2 Relation between patient characteristics and mortality, separately examining deaths from natural causes only and deaths that combine natural causes with accident and misadventure verdicts.

	Death from natural causes		Death from natural cause or accident/misadventure	
	Proportion who died if risk factor present; not present (values are numbers of deaths/people at risk)	Difference between proportions (95% confidence interval)	Proportion who died if risk factor present; not present (values are numbers of deaths/people at risk)	Difference between proportions (95% confidence interval)
Characteristics pre-dating index episode:				
Serious illness	29/87 (33%); 37/493 (8%)	25.8% (16.4 to 36.4)	32/90 (36%); 46/502 (9%)	26.4% (16.9 to 36.9)
Live alone	26/119 (22%); 49/538 (9%)	12.7% (5.7 to 21.3)	31/124 (25%); 59/548 (11%)	14.2% (6.9 to 22.9)
Previous self-harm	40/236 (17%); 24/380 (6%)	10.6% (5.5 to 16.3)	48/244 (20%); 30/386 (8%)	11.9% (6.4 to 17.8)
Psychiatric history	41/226 (18%); 24/367 (7%)	11.6% (6.2 to 17.5)	50/235 (21%); 29/372 (8%)	13.5% (7.8 to 19.6)
Previous psychiatric admission	30/122 (25%); 31/452 (7%)	17.7% (10.4 to 26.3)	33/125 (26%); 40/461 (9%)	17.7% (10.2 to 26.4)
Characteristics of the episode				
Attend during day (0800-1600)	43/228 (19%); 58/624 (9%)	9.6% (4.4 to 15.5)	52/237 (22%); 72/638 (11%)	10.7% (5.2 to 16.8)
Drowsy rather than alert	68/387 (18%); 34/470 (7%)	10.3% (5.9 to 14.9)	81/400 (20%); 44/480 (9%)	11.1% (6.4 to 15.9)
Poisoned with psychotropic drugs or non-ingestibles	74/375 (20%); 28/482 (6%)	13.9% (9.5 to 18.6)	88/389 (23%); 37/491 (8%)	15.1% (10.4 to 19.9)
Subsequent episodes of self-harm				
Repeated within 1 year of index episode	18/100 (18%); 84/757 (11%)	6.9% (0.15 to 15.8)	22/104 (21%); 103/776 (13%)	7.9% (0.67 to 17.0)

1
2
3
4 **ACKNOWLEDGEMENTS**
5

6 We thank the Office for National Statistics for provision of data. Tracing of the
7
8 patients by the Office for National Statistics was supported by a research grant
9
10 from the former NHS Northern and Yorkshire Region.
11
12
13
14

15 **Conflict of interest statement** None
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

REFERENCES

1. Antretter E, Dunkel D, Haring C (2009) Cause-specific excess mortality in suicidal patients: gender differences in mortality patterns. *Gen Hosp Psychiat* 31: 67-74
2. Carter G, Reith DM, Whyte IM, McPherson M (2005) Non-suicidal deaths following hospital-treated self-poisoning. *Aust NZ J Psychiat* 39: 101-107
3. Chang SS, Sterne JAC, Lu T-H, Gunnell D. (2010) 'Hidden' suicides amongst deaths certified as undetermined intent, accident by pesticide poisoning and accident by suffocation in Taiwan. *Soc Psychiat Epidemiol* 45: 143-152
4. Charlton J, Kelly S, Dunnell K, Evans B, Jenkins R, Wallis R (1992) Trends in suicide deaths in England and Wales. *Population Trends* 69: 10-16
5. De Leo D, Scocco P, Marietta P, Schmidtke A, Bille-Brahe U, Kerkhof AJFM, Lönnqvist J, Crepet P, Salander-Renberg E, Wasserman D, Michel K, Bjerke T (1999) Physical illness and parasuicide: evidence from the European Parasuicide Study Interview Schedule (EPSIS/ WHO-EURO). *Int J Psychiat Med* 29: 149-163
6. Dennis M, Read S, Andrews H, Wakefield P, Zafar R, Kavi S (2001) Suicide in a single health district: epidemiology and audit of the involvement of psychiatric services. *J Ment Health* 6: 673-682

7. Dennis M, Baillon S, Brugha T, Lindesay J, Stewart R, Meltzer H (2007)
The spectrum of suicidal ideation in Great Britain: comparisons across a
16-74 age range. *Psychol Med* 37: 795-805
8. Ekeberg O, Ellingsen O, Jacobsen D (1991) Suicide and other causes of
death in a five-year follow-up of patients treated for self-poisoning in Oslo.
Acta Psychiat Scand 83: 432–437
9. Ekeberg O, Ellingsen O, Jacobsen D (1994) Mortality and causes of death
in a 10-year follow-up of patients treated for self-poisonings in Oslo.
Suicide Life-Threat 24: 398-405
10. Hall DJ, O'Brien F, Stark C, Pelosi A, Smith H (1998) Thirteen-year follow-
up of deliberate self-harm, using linked data. *Brit J Psychiat* 172: 239-242
11. Harris EC, Barraclough BM (1994) Suicide as an outcome for medical
disorders. *Medicine* 73: 281-296
12. Hawton K, Townsend E, Arensman E, Gunnell D, Hazell P, House A, van
Heeringen K (1999) Psychosocial and pharmacological treatments for
deliberate self harm. *Cochrane Db Syst Rev*. Issue 4. Art. No.: CD001764.
DOI: 10.1002/14651858.CD001764

13. Hawton K, Bergen H, Casey D, Simkin S, Palmer B, Cooper J, Kapur N, Horrocks J, House A, Lilley R, Noble R, Owens D (2007) Self-harm in England: a tale of three cities. Soc Psych Psych Epid 42: 513-521
14. Hawton K, Harriss L (2007) Deliberate self-harm in young people: characteristics and subsequent mortality in a 20-year cohort of patients presenting to hospital. J Clin Psychiat 68: 1574-1583
15. Hawton K, Harriss L, Hall S, Simkin S, Bale E, Bond, A (2003) Deliberate self-harm in Oxford, 1990-2000: a time of change in patient characteristics. Psychol Med 33: 987-996
16. Hawton K, Harriss L, Zahl, D (2006) Death from all causes in a long-term follow-up study of 11 583 deliberate self-harm patients. Psychol Med 36: 397-405
17. Holley HL, Fick G, Love EJ (1998) Suicide following an inpatient hospitalization for a suicide attempt. Soc Psych Psych Epid 33: 543-551
18. Jenkins GR, Hale R, Papanastassiou M, Crawford MJ, Tyrer P (2002) Suicide rate 22 years after parasuicide: cohort study. BMJ 325: 1155
19. Neeleman J (2001) A continuum of premature death. Meta-analysis of competing mortality in the psychologically vulnerable. Int J Epidemiol 30: 154-162

- 1
2
3
4 20. NICE (National Institute for Clinical Excellence) (2004) Short-term physical
5 and psychological management and secondary prevention of self-harm in
6 primary and secondary care. NICE, London
7
8
9
10
11
12
13 21. Nordentoft M, Breum L, Munck LK, Nordestgaard AG, Hunding A, Laursen
14 Bjaeldager PA (1993) High mortality by natural and unnatural causes: a 10
15 year follow up study of patients admitted to a poisoning treatment centre
16 after suicide attempt. BMJ 306: 1637-1641
17
18
19
20
21
22
23
24 22. Ostamo A, Lönnqvist J (2001) Excess mortality of suicide attempters. Soc
25 Psych Psych Epid 36: 29-35
26
27
28
29
30
31 23. Owens D, Horrocks J, House A (2002) Fatal and non-fatal repetition of
32 self-harm: a systematic review. Br J Psychiat 181: 193-199
33
34
35
36
37
38 24. Owens D, Dennis M, Jones S, Dove A, Dave S (1991) Self-poisoning
39 patients discharged from accident and emergency: risk factors and
40 outcome. J Roy Coll Phys Lond 25: 218-222
41
42
43
44
45
46 25. Owens D, Wood C, Greenwood D, Hughes T, Dennis M (2005) Mortality
47 and suicide after non-fatal self-poisoning: a 16-year outcome study of
48 patients attending accident and emergency. Br J Psychiat 187: 470-475
49
50
51
52
53
54
55 26. Rygnestad, T. (1997) Mortality after deliberate self-poisoning. A
56 prospective follow-up study of 587 persons observed for 5279 person
57
58
59
60
61
62
63
64
65

years: risk factors and causes of death. Soc Psych Psych Epid 32: 443-450.

27. Sakinofsky I (2000) Repetition of suicidal behaviour. In: Hawton K, van Heeringen K (eds) The international handbook of suicide and attempted suicide. Wiley, Chichester, pp 385-404
28. Sinclair JMA, Gray A, Rivero-Arias O, Saunders KEA, Hawton K. (2010) Healthcare and social services resource use and costs of self-harm patients. Soc Psychiat Epidemiol DOI 10.1007/s00127-010-0183-5
29. Start R, Delargy-Aziz Y, Dorries C, Silcocks P, Cotton D. (1993) Clinicians and the coronial system: Ability of clinicians to recognise reportable deaths. BMJ 306: 1038-1041
30. Stenager EN, Stenager E (2000) Physical illness and suicidal behaviour. In: Hawton K, van Heeringen K (eds) The international handbook of suicide and attempted suicide. Wiley, Chichester, pp 405-420
31. Suominen K, Isomestä E, Haukka J, Lönnqvist J (2004) Substance use and male gender as risk factors for deaths and suicide. A 5-year follow-up study after deliberate self-harm. Soc Psych Psych Epid 39: 720-724
32. World Health Organization (1993) The ICD-10 classification of mental and behavioural disorders: diagnostic criteria for research. WHO, Geneva